

Microdermabrasion Kingston

Microdermabrasion Kingston - The condition of Dermatitis Herpetiformis is a chronic skin condition characterized by blisters filled with a watery fluid. It is likewise referred to as DH or Duhring's Disease. The name implies the meaning in that it is a skin disease which has the specific appearance like herpes, however, the condition is not caused by or related to the herpes virus. DH was initially described by Dr. Louis Duhring during 1884, thus the name. In the year 1867, a correlation between celiac disease or gluten intolerance and DH was recognized. In most of cases, the age of onset is normally 15 to 40 years, even if, DH could likewise affect children as well as the elderly. Women and men are equally affected. The prevalence of DH varies from 1 in 400 to 1 in 10,000.

Signs

Dermatitis Herpetiformis has a characteristic rash that contains chronic papulovesicular eruptions, which are intensely itchy. This particular kind of rash is normally distributed symmetrically on extensor surfaces like the back of neck, elbows, knees, buttocks and the back. The blisters could vary from being up to 1 cm across to very small. Because this particular condition is very itchy intensive, the need to scratch can be overwhelming. This may cause the individual to really scratch blisters off before they could be examined by a doctor of medicine. Intense itching or intense burning sensation are sometimes felt prior to blisters appearing in a particular part.

The severity of Dermatitis Herpetiformis could differ in time if left untreated. Quite often, the first symptoms appear in early adulthood, between the ages of 20 and 30 years old. The very first visible signs are the tiny papules or vesicles which resemble blisters or bumps. The initial symptoms are intensive burning and itching. Usually, the first blisters appear on the shoulders, on the lower end of the spinal column, along the hairline and at times in the mouth. Rarely, does the rash happen on several mucous membranes other than the lips or the mouth. If gluten ingestion is avoided and the right treatment is given, the signs normally disappear. In some cases, taking oral contraceptives could worsen the symptoms.

The symptoms of DH tend to come and go, in short periods of time, yet are always persistent. These symptoms can be accompanied by fatigue and abdominal pain and some signs of celiac disease.

The DH rash typically forms and disappears in 3 stages. The person would at first see a slight discoloration of the skin where the lesions appear. The next stage will be the lesions transforming into groups of obvious papules and vesicles. The healing stage of the lesions is the last stage of development. Often this set of indications is characterized by a skin color change. This might lead to areas of the skin turning darker or lighter than the color of the skin somewhere else on the body. Because of the intense itching, individuals generally scratch that can lead to the formation of crusts.